



ICLUSIG[®]

(ponatinib) tablets
45mg / 30mg / 15mg / 10mg

Access Guide



INDICATIONS AND USAGE

ICLUSIG is indicated for the treatment of adult patients with:

Philadelphia Chromosome-Positive Acute Lymphoblastic Leukemia (Ph+ ALL)

- Newly diagnosed Ph+ ALL in combination with chemotherapy.
This indication is approved under accelerated approval based on minimal residual disease (MRD)-negative complete remission (CR) at the end of induction. Continued approval for this indication may be contingent upon verification of clinical benefit in a confirmatory trial(s).
- As monotherapy in Ph+ ALL for whom no other kinase inhibitors are indicated or T315I-positive Ph+ ALL.

Chronic Myeloid Leukemia (CML)

- Chronic phase (CP) CML with resistance or intolerance to at least two prior kinase inhibitors.
- Accelerated phase (AP) or blast phase (BP) CML for whom no other kinase inhibitors are indicated.
- T315I-positive CML (chronic phase, accelerated phase, or blast phase).

Limitations of Use: ICLUSIG is not indicated and is not recommended for the treatment of patients with newly diagnosed CP-CML.

IMPORTANT SAFETY INFORMATION

WARNING: ARTERIAL OCCLUSIVE EVENTS, VENOUS THROMBOEMBOLIC EVENTS, HEART FAILURE, and HEPATOTOXICITY

See full prescribing information for complete boxed warning.

- Arterial occlusive events (AOEs), including fatalities, have occurred in ICLUSIG-treated patients. AOEs included fatal myocardial infarction, stroke, stenosis of large arterial vessels of the brain, severe peripheral vascular disease, and the need for urgent revascularization procedures. Patients with and without cardiovascular risk factors, including patients age 50 years or younger, experienced these events. Monitor for evidence of AOEs. Interrupt or discontinue ICLUSIG based on severity. Consider benefit-risk to guide a decision to restart ICLUSIG.
- Venous thromboembolic events (VTEs) have occurred in ICLUSIG-treated patients. Monitor for evidence of VTEs. Interrupt or discontinue ICLUSIG based on severity.
- Heart failure, including fatalities, occurred in ICLUSIG-treated patients. Monitor for heart failure and manage patients as clinically indicated. Interrupt or discontinue ICLUSIG for new or worsening heart failure.
- Hepatotoxicity, liver failure and death have occurred in ICLUSIG-treated patients. Monitor liver function tests. Interrupt or discontinue ICLUSIG based on severity.

Please see [full Prescribing Information](#), including Boxed Warning and Important Safety Information throughout.



Please see [page 3](#) to learn more.



ONCOLOGY

Distribution Details for ICLUSIG® (ponatinib)

Product name ICLUSIG® (ponatinib) tablets, for oral use

Distributed by Takeda Pharmaceuticals America, Inc.

Storage and handling Store ICLUSIG tablets at 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F)

Recommended dosing

Newly diagnosed Ph+ ALL: The recommended starting dosage of ICLUSIG in combination with chemotherapy is 30 mg orally once daily with a reduction to 15 mg orally once daily upon achievement of MRD-negative ($\leq 0.01\%$ BCR::ABL1/ABL1) complete remission (CR) at the end of induction. Continue ICLUSIG in combination with chemotherapy for up to 20 cycles until loss of response or unacceptable toxicity.

Monotherapy for Ph+ ALL for whom no other kinase inhibitors are indicated or T3151-positive Ph+ ALL: The optimal dose of ICLUSIG has not been identified. The recommended starting dosage of ICLUSIG is 45 mg orally once daily. Continue ICLUSIG until loss of response or unacceptable toxicity. Consider discontinuing ICLUSIG if response has not occurred by 3 months.

CP-CML: The recommended starting dosage of ICLUSIG is 45 mg orally once daily with a reduction to 15 mg orally once daily upon achievement of $\leq 1\%$ BCR::ABL1[®]. Patients with loss of response can re-escalate the dose of ICLUSIG to a previously tolerated dosage of 30 mg or 45 mg orally once daily. Continue ICLUSIG until loss of response at the re-escalated dose or unacceptable toxicity. Consider discontinuing ICLUSIG if hematologic response has not occurred by 3 months.

AP-CML and BP-CML: The optimal dose of ICLUSIG has not been identified. The recommended starting dosage of ICLUSIG is 45 mg orally once daily. Consider reducing the dose of ICLUSIG for patients with AP-CML who have achieved a major cytogenetic response. Continue ICLUSIG until loss of response or unacceptable toxicity. Consider discontinuing ICLUSIG if response has not occurred by 3 months.

How supplied	Strength	Count	NDC Number
	45 mg	Bottle of 30 tablets	63020-534-30
	30 mg	Bottle of 30 tablets	63020-533-30
	15 mg	Bottle of 30 tablets	63020-535-30
	10 mg	Bottle of 30 tablets	63020-536-30

ICLUSIG tablets are supplied in wide-mouth white high-density polyethylene (HDPE) bottles with a desiccant canister and induction-sealed, child-resistant closure.

Ordering

ICLUSIG is available through an exclusive, integrated specialty pharmacy:
AcariaHealth
 833-442-8911
www.acariahealth.com

There are 2 ways to order ICLUSIG via the AcariaHealth Specialty Pharmacy:

1. Submit the ICLUSIG prescription to any AcariaHealth in your e-prescribing system
2. Complete the ICLUSIG Referral Form provided by AcariaHealth at acariahealth.envolvehealth.com/resources/referral-forms.html and fax to AcariaHealth as directed on the form

ICLUSIG must be filled through AcariaHealth. Sending an ICLUSIG prescription to an alternate pharmacy may result in delay or nonfulfillment of the prescription.

ICLUSIG may be purchased directly from our distribution partner:
 Foundation Care,
 An AcariaHealth Solution
 ICLUSIGDirect Purchase Program
 833-291-2773
www.ICLUSIGDirect.com

AP, accelerated phase; BCR::ABL1, breakpoint cluster region-Abelson; BP, blast phase; CP, chronic phase; CML, chronic myeloid leukemia; MRD, minimal residual disease; NDC, National Drug Code; Ph+ ALL, Philadelphia chromosome positive acute lymphoblastic leukemia; USP, United States Pharmacopeia.



Please see [full Prescribing Information](#), including Boxed Warning and Important Safety Information throughout.



We're here to help your patients with their coverage, financial, and educational resource needs

Committed to supporting your patients

Takeda Oncology Here2Assist® is a comprehensive support program committed to helping your patients navigate coverage requirements, identify available financial assistance, and connect with helpful resources throughout their Takeda Oncology treatment.

- ▶ Works with your patients' insurance company to help get your patient started on their medication
- ▶ Identifies available financial assistance that may be right for your patients
- ▶ Identifies specialty pharmacies to help fill and ship your patients' prescriptions appropriately
- ▶ Conducts regular follow-up calls to patients

Visit us at <https://www.here2assist.com/hcp/home> to learn more

Daryl
Takeda Oncology
Here2Assist® patient



IMPORTANT SAFETY INFORMATION (CONT'D)

WARNINGS AND PRECAUTIONS

Arterial Occlusive Events (AOEs): AOE, including fatalities, have occurred in patients who received ICLUSIG in PhALLCON, OPTIC and PACE. These included cardiovascular, cerebrovascular, and peripheral vascular events. In PhALLCON, 6% of 163 patients experienced AOE; 3.7% experienced Grade 3 or 4. The incidence of AOE in OPTIC (45 mg→15 mg) was 14% of 94 patients; 6% experienced Grade 3 or 4. In PACE, the incidence of AOE was 26% of 449 patients; 14% experienced Grade 3 or 4. Fatal AOE occurred in 0.6% of patients in PhALLCON, 2.1% of patients in OPTIC, and in 2% of patients in PACE. Some patients in PACE experienced recurrent or multisite vascular occlusion. Patients with and without cardiovascular risk factors, including patients age 50 years or younger, experienced these events. The most common risk factors observed with these events in PACE were history of hypertension, hypercholesterolemia, and non-ischemic cardiac disease. In PhALLCON, OPTIC and PACE, AOE were more frequent with increasing age.

In PhALLCON, patients with uncontrolled hypertension, hypertriglyceridemia, or diabetes were excluded. Patients with clinically significant, uncontrolled, or active cardiovascular disease, including any history of myocardial infarction, peripheral vascular infarction, revascularization procedure, venous thromboembolism, clinically significant atrial/ventricular tachyarrhythmias, unstable angina, or congestive heart failure within the 6 months prior to the first dose of ICLUSIG, were also excluded.

In OPTIC, patients with uncontrolled hypertension or diabetes and patients with clinically significant, uncontrolled, or active cardiovascular disease were excluded.


In PACE, patients with uncontrolled hypertriglyceridemia and patients with clinically significant or active cardiovascular disease within the 3 months prior to the first dose of ICLUSIG were excluded.

Consider whether the benefits of ICLUSIG are expected to exceed the risks. Monitor for evidence of AOE. Interrupt, then resume at the same or decreased dose or discontinue ICLUSIG based on recurrence/severity. Consider benefit-risk to guide a decision to restart ICLUSIG.

Venous Thromboembolic Events (VTEs): Serious or severe VTEs have occurred in patients who received ICLUSIG. In PhALLCON, VTEs occurred in 12% of 163 patients, including serious or severe (Grade 3 or 4) in 3.1% of patients. One of 94 patients in OPTIC experienced a VTE (Grade 1 retinal vein occlusion). In PACE, VTEs occurred in 6% of 449 patients including serious or severe (Grade 3 or 4) VTEs in 5.8% of patients. In PhALLCON and PACE VTEs included deep venous thrombosis, embolism, pulmonary embolism, superficial vein thrombosis, thrombosis, jugular vein thrombosis, superficial thrombophlebitis, retinal vein occlusion, and retinal vein thrombosis with vision loss. The incidence of VTEs in PACE was higher in patients with Ph+ ALL (9% of 32 patients) and BP-CML (10% of 62 patients). Monitor for evidence of VTEs. Interrupt, then resume at the same or decreased dose or discontinue ICLUSIG based on recurrence/severity.

Heart Failure: Fatal, serious or severe heart failure events have occurred in patients who received ICLUSIG. In PhALLCON, heart failure occurred in 6% of 163 patients; 1.2% experienced serious or severe (Grade 3 or 4) heart failure. Heart failure occurred in 13% of 94 patients in OPTIC; 1.1% experienced serious or severe (Grade 3 or 4). In PACE, heart failure occurred in 9% of 449 patients; 7% experienced serious or severe (Grade 3 or higher). In PhALLCON the most frequently reported heart failure event (>1 patient) was increased brain natriuretic peptide (BNP) (2.5%). In OPTIC, the most frequently reported heart failure events (>1 patient each) were left ventricular hypertrophy (3.2%) and BNP increased (3.2%). In PACE, the most frequently reported heart failure events (≥2%) were congestive cardiac failure (3.1%), decreased ejection fraction (2.9%), and cardiac failure (2%). Monitor patients for signs or symptoms consistent with heart failure and manage heart failure as clinically indicated. Interrupt, then resume at reduced dose or discontinue ICLUSIG for new or worsening heart failure. Monitor patients for signs or symptoms consistent with heart failure and manage heart failure as clinically indicated. Interrupt, then resume at reduced dose or discontinue ICLUSIG for new or worsening heart failure.

Please see [full Prescribing Information](#), including Boxed Warning and Important Safety Information throughout.



IMPORTANT SAFETY INFORMATION (CONT'D)

WARNINGS AND PRECAUTIONS (CONT'D)

Hepatotoxicity: ICLUSIG can cause hepatotoxicity, including liver failure and death. Fulminant hepatic failure leading to death occurred in 3 patients, with hepatic failure occurring within 1 week of starting ICLUSIG in one of these patients. These fatal cases occurred in patients with BP-CML or Ph+ ALL treated with monotherapy. Hepatotoxicity occurred in 66% of 163 patients in PhALLCON, in 28% of 94 patients in OPTIC and in 32% of 449 patients in PACE. Grade 3 or 4 hepatotoxicity occurred in PhALLCON (30% of 163 patients), in OPTIC (6% of 94 patients), and in PACE (13% of 449 patients). The most frequent hepatotoxic events were elevations of ALT, AST, GGT, bilirubin, and alkaline phosphatase. Monitor liver function tests at baseline, then at least monthly or as clinically indicated. Interrupt, then resume at a reduced dose or discontinue ICLUSIG based on recurrence/severity.

Hypertension: Serious or severe hypertension, including hypertensive crisis, has occurred in patients who received ICLUSIG. Patients may require urgent clinical intervention for hypertension associated with confusion, headache, chest pain, or shortness of breath. Monitor blood pressure at baseline and as clinically indicated and manage hypertension as clinically indicated. Interrupt, dose reduce, or stop ICLUSIG if hypertension is not medically controlled. For significant worsening, labile or treatment-resistant hypertension, interrupt ICLUSIG and consider evaluating for renal artery stenosis.

Pancreatitis: Serious or severe pancreatitis has occurred in patients who received ICLUSIG. Elevations of lipase and amylase also occurred. In the majority of cases that led to dose modification or treatment discontinuation, pancreatitis resolved within 2-3 weeks. Monitor serum lipase every 2 weeks for the first 2 months and then monthly thereafter or as clinically indicated. Consider additional serum lipase monitoring in patients with a history of pancreatitis or alcohol abuse. Interrupt, then resume at the same or reduced dose or discontinue ICLUSIG based on severity. Evaluate for pancreatitis when lipase elevation is accompanied by abdominal symptoms.

Increased Toxicity in Newly Diagnosed Chronic Phase CML: In a prospective randomized clinical trial in the first-line treatment of newly diagnosed patients with CP-CML, single agent ICLUSIG 45 mg once daily increased the risk of serious adverse reactions 2-fold compared to single agent imatinib 400 mg once daily. The median exposure to treatment was less than 6 months. The trial was halted for safety. Arterial and venous thrombosis and occlusions occurred at least twice as frequently in the ICLUSIG arm compared to the imatinib arm. Compared to imatinib-treated patients, ICLUSIG-treated patients exhibited a greater incidence of myelosuppression, pancreatitis, hepatotoxicity, cardiac failure, hypertension, and skin and subcutaneous tissue disorders. ICLUSIG is not indicated and is not recommended for the treatment of patients with newly diagnosed CP-CML.

Neuropathy: Peripheral and cranial neuropathy occurred in patients in PhALLCON, OPTIC and PACE. Some of these events in PhALLCON and PACE were Grade 3 or 4. Monitor patients for symptoms of neuropathy, such as hypoesthesia, hyperesthesia, paresthesia, discomfort, a burning sensation, neuropathic pain or weakness. Interrupt, then resume at the same or reduced dose or discontinue ICLUSIG based on recurrence/severity.

Ocular Toxicity: Serious or severe ocular toxicity leading to blindness or blurred vision have occurred in ICLUSIG-treated patients. The most frequent ocular toxicities occurring in PhALLCON, OPTIC and PACE were dry eye, blurred vision, and eye pain. Retinal toxicities included age-related macular degeneration, macular edema, retinal vein occlusion, retinal hemorrhage, and vitreous floaters. Conduct comprehensive eye exams at baseline and periodically during treatment.

Please see [full Prescribing Information](#), including Boxed Warning and Important Safety Information throughout.

 **ICLUSIG**[®]
(ponatinib) tablets
45mg / 30mg / 15mg / 10mg

IMPORTANT SAFETY INFORMATION (CONT'D)

WARNINGS AND PRECAUTIONS (CONT'D)

Hemorrhage: Fatal and serious hemorrhage events have occurred in patients who received ICLUSIG. Fatal hemorrhages occurred in PACE and serious hemorrhages occurred in PhALLCON, OPTIC and PACE. In PACE, the incidence of serious bleeding events was higher in patients with AP-CML, BP-CML, and Ph+ ALL. Intracranial hemorrhage, gastrointestinal hemorrhage and subdural hematoma were the most frequently reported serious hemorrhages. Events often occurred in patients with Grade 4 thrombocytopenia. Monitor for hemorrhage and manage patients as clinically indicated. Interrupt, then resume at the same or reduced dose or discontinue ICLUSIG based on recurrence/severity.

Fluid Retention: Fatal and serious fluid retention events have occurred in patients who received ICLUSIG. In PACE, one instance of brain edema was fatal and serious events included pleural effusion, pericardial effusion, and angioedema. In PhALLCON serious fluid retention included pericardial effusion. The most frequent occurrences of fluid retention in patients who received ICLUSIG were peripheral edema and pleural effusion. Monitor for fluid retention and manage patients as clinically indicated. Interrupt, then resume at the same or reduced dose or discontinue ICLUSIG based on recurrence/severity.

Cardiac Arrhythmias: Cardiac arrhythmias, including ventricular, atrial arrhythmias, tachycardia, syncope, atrial fibrillation and supraventricular tachycardia occurred in patients in PhALLCON, OPTIC, and PACE. For some patients, events were serious or severe (Grade 3 or 4) and led to hospitalization. Monitor for signs and symptoms suggestive of slow heart rate (fainting, dizziness) or rapid heart rate (chest pain, palpitations or dizziness) and manage patients as clinically indicated. Interrupt, then resume at the same or reduced dose or discontinue ICLUSIG based on recurrence/severity.

Myelosuppression: Grade 3 or 4 events of neutropenia, thrombocytopenia, and anemia occurred in patients in PhALLCON, OPTIC and PACE. In PACE, the incidence of myelosuppression was greater in patients with AP-CML, BP-CML, and Ph+ ALL treated with monotherapy than in patients with CP-CML. Obtain complete blood counts every 2 weeks for the first 3 months and then monthly or as clinically indicated. If ANC less than $1 \times 10^9/L$ or platelets less than $50 \times 10^9/L$, interrupt ICLUSIG until ANC at least $1.5 \times 10^9/L$ and platelets at least $75 \times 10^9/L$, then resume at same or reduced dose.

Tumor Lysis Syndrome (TLS): Serious TLS was reported in ICLUSIG-treated patients in PhALLCON, OPTIC and PACE. Ensure adequate hydration and treat high uric acid levels prior to initiating ICLUSIG.

Reversible Posterior Leukoencephalopathy Syndrome (RPLS): RPLS (also known as Posterior Reversible Encephalopathy Syndrome) has been reported in patients who received ICLUSIG. Patients may present with neurological signs and symptoms, visual disturbances, and hypertension. Diagnosis is made with supportive findings on magnetic resonance imaging (MRI) of the brain. Interrupt ICLUSIG until resolution. The safety of resumption of ICLUSIG in patients upon resolution of RPLS is unknown.

Impaired Wound Healing and Gastrointestinal Perforation: Impaired wound healing occurred in patients receiving ICLUSIG. Withhold ICLUSIG for at least 1 week prior to elective surgery. Do not administer for at least 2 weeks following major surgery and until adequate wound healing. The safety of resumption of ICLUSIG after resolution of wound healing complications has not been established. Gastrointestinal perforation or fistula occurred in patients receiving ICLUSIG. Permanently discontinue in patients with gastrointestinal perforation.

Embryo-Fetal Toxicity: Based on its mechanism of action and findings from animal studies, ICLUSIG can cause fetal harm when administered to a pregnant woman. Advise pregnant women of the potential risk to the fetus. Advise females of reproductive potential to use effective contraception during treatment with ICLUSIG and for 3 weeks after the last dose.

Please see [full Prescribing Information](#), including Boxed Warning and Important Safety Information throughout.

 **ICLUSIG**[®]
(ponatinib) tablets
45mg / 30mg / 15mg / 10mg

IMPORTANT SAFETY INFORMATION (CONT'D)

ADVERSE REACTIONS

The most common adverse reactions (occurring in >20% of patients) are:

- ICLUSIG as a single agent: rash and related conditions, arthralgia, abdominal pain, headache, constipation, dry skin, hypertension, fatigue, fluid retention and edema, pyrexia, nausea, pancreatitis/lipase elevation, hemorrhage, anemia, hepatic dysfunction and AOE. The most common Grade 3 or 4 laboratory abnormalities (>20%) are platelet count decreased, neutrophil cell count decreased, and white blood cell decreased.
- ICLUSIG in combination with chemotherapy: hepatic dysfunction, arthralgia, rash and related conditions, headache, pyrexia, abdominal pain, constipation, fatigue, nausea, oral mucositis, hypertension, pancreatitis/lipase elevation, neuropathy peripheral, hemorrhage, febrile neutropenia, fluid retention and edema, vomiting, paresthesia and cardiac arrhythmias. The most common Grade 3 or 4 laboratory abnormalities (>20%) are decreased white blood cell count, decreased neutrophil cell count, decreased platelet count, decreased lymphocyte cell count, decreased hemoglobin, increased lipase and increased alanine aminotransferase.

To report SUSPECTED ADVERSE REACTIONS, contact Takeda Pharmaceuticals at 1-844-817-6468 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Strong CYP3A Inhibitors: Avoid coadministration or reduce ICLUSIG dose if coadministration cannot be avoided.

Strong CYP3A Inducers: Avoid coadministration.

USE IN SPECIFIC POPULATIONS

Lactation: Advise women not to breastfeed during treatment with ICLUSIG and for 1 week following last dose.

Females and Males of Reproductive Potential: Verify pregnancy status of females of reproductive potential prior to initiating ICLUSIG. Ponatinib may impair fertility in females, and it is not known if these effects are reversible.

Pre-existing Hepatic Impairment: For patients with CP-CML, AP-CML, BP-CML, and Ph+ ALL receiving monotherapy, reduce the starting dose of ICLUSIG to 30mg orally once daily for patients with pre-existing hepatic impairment as these patients are more likely to experience adverse reactions compared to patients with normal hepatic function. For patients with newly diagnosed Ph+ ALL, no dosage adjustment is recommended.

Please see accompanying [full Prescribing Information](#), including **Boxed Warning**.

To learn more about ICLUSIG, please visit www.iclusig.com/hcp.

Reference: ICLUSIG. Prescribing information. Takeda Pharmaceuticals U.S.A., Inc.; March 2024.



ONCOLOGY



ICLUSIG[®]
(ponatinib) tablets
45mg / 30mg / 15mg / 10mg

TAKEDA[®] and the TAKEDA Logo[®] are registered trademarks of Takeda Pharmaceutical Company Limited.
ICLUSIG[®] and the ICLUSIG Logo[®] are registered trademarks of ARIAD Pharmaceuticals, Inc.
HERE2ASSIST[®] and the HERE2ASSIST Logo[®] are registered trademarks of Millennium Pharmaceuticals, Inc.
© 2024 Takeda Pharmaceuticals U.S.A., Inc. All rights reserved. 03/24 USO-ICL-0562